

Amendment-1 Annexure-5, RMP Grapes 2011 season

Apeda/Q/Grapenet/2010-11/

Date: 10.09.2010

To all stakeholders for export of fresh table grapes

Sub: RMP Grapes for 2011 season amendment in Annexure-5 by NRC Grapes Pune

Dear Sirs,

Please refer to our email dated 20.08.2010 circulating RMP for grapes for 2011 season. Subsequently NRC Grapes Pune has modified Annexure-5 of the RMP removing the following chemicals from the list as they do not have label claim for usage in grapes:

1. Copper hydroxide 77WP
2. Bordeaux mixture
3. Potassium bicarbonate
4. Azadiechtin (0.03%)

The modified Annexure-5 is enclosed as submitted by NRC Grapes Pune dated 04 September, 2010 replacing the Annexure-5 of RMP Grapes 2011 dated 18.08.2010 circulated on 20.08.2010 to all. You may please ensure necessary compliances by the respective farmers, State Governments and other stakeholders.

With regards,

S. Dave  
Director - APEDA

Modification date: 04 September, 2010

## List of chemicals with CIB&amp;RC label claim for use in grapes



NATIONAL RESEARCH CENTRE FOR GRAPES  
(Indian Council of Agricultural Research)

P.B. No. 3, Manjri Farm P.O. Solapur Road Pune - 412 307, India

Tel.: 020-26914245, 25169101 Fax: 020-26914246

E-mail: nrcgrapes@gmail.com, Website: <http://nrcgrapes.mah.nic.in>

Sr. No.	Chemical recommended for major disease & pest	Nature of chemical	Dose on formulation basis	EU MRL (mg/kg) updated as on 19/07/2010	Pre-harvest Interval (PHI in days)
I	Downy Mildew				
1.	Mancozeb 75 WP	NS	1.5-2.0 g/L	5.0	35 (avoid using after fruit set)
2.	Propineb 70 WP	NS	3.0 g/L	1.0	40 (avoid using after fruit set)
3.	COC 50 WP	NS	2.5 g/L, 2.4 g/L	50.0	42 (avoid using after fruit set)
4.	Chlorothalonil 75 WP	NS	2.0 g/L	1.0	60
5.	Fosetyl AL 80 WP	S	1.4-2.0 g/L	100.0	7
6.	Metalaxyl + Mancozeb 8+64 WP	S+NS	2.5 g/L	2.0 + 5.0	66
6a.	Metalaxyl-M + Mancozeb 4+64 WP	S+NS	2.5 g/L	2.0 + 5.0	66
7.	Cymoxanil + Mancozeb 8+64 WP	S+NS	2.0 g/L	0.2 + 5.0	66
8.	Dimethomorph 50 WP + Mancozeb 75WP as tank mixture	S+NS	0.5 to 0.75 g/L + 2.0 g/L	3.0 + 5.0	66
9.	Fenamidone + Mancozeb 10+50 WG	S+NS	2.5 to 3 g/L	0.5 + 5.0	66
10.	Azoxystrobin 23 SC	S	200 mL/Acre	2.0	7
11.	Iprovalicarb + Propineb 5.5+61.25WP	S+NS	2.25 g/L	2.0 + 1.0	55
12.	Famoxadone 16.6 % + Cymoxanil 22.1 % SC	S+NS	500 mL/ha	2.0 + 0.2	27
13.	Kresoxim methyl 44.3 SC	S	600-700 mL/ha	1.0	30
14.	Fenamidone 4.44% + Fosetyl-Al 66.66% WDG	S	2 to 2.5 kg/ha	0.5 + 100	27
II	Powdery Mildew				

Sr. No.	Chemical recommended for major disease & pest	Nature of chemical	Dose on formulation basis	EU MRL (mg/kg) updated as on 19/07/2010	Pre-harvest Interval (PHI in days)
15.	Penconazole 10 EC	S	0.50 mL/L	0.2	50
16.	Triadimefon 25 WP	S	0.50-1.0 g/L	2.0	45
17.	Hexaconazole 5EC	S	1.0 mL/L	0.1	38
18.	Myclobutanil 10 WP	S	0.40 g/L	1.0	30
19.	Flusilazole 40 EC	S	25 mL / 200 L	0.05	50
20.	Fenarimol 10 EC	S	0.40 mL / L	0.3	30
21.	Difenoconazole 25EC	S	0.50 mL / L	0.5	45
10a.	Azoxystrobin 23 SC	S	200 mL / Acre	2.0	7
13a.	Kresoxim methyl 44.3 SC	S	600-700 mL/ha	1.0	30
22.	Dinocap 48 EC	NS	0.30 - 0.35 mL/L	0.05	50 (avoid application when tender shoots are present in canopy)
23.	Sulfur 40 SC, 55.16 SC, 80 WP, 80 WDG, 85 WP	NS	3.0 mL, 3.0 mL, 2.50 g, 1.87-2.50 g, 1.50-2.0 g/L, respectively	MRL not applicable	--
III	Anthracnose				
2a.	Propineb 70 WP	NS	3.0 g/L	1.0	40
3a.	COC 50 WP	NS	2.5 g/L, 2.40 g/L	50.0	42 (avoid using after fruit set)
24.	Carbendazim 50 WP, 46.27 SC	S	1.0 g/L, 1.0 mL/L	0.3	50
IV	Post harvest berry rots				
	--				
V	Flea beetles				
25.	Imidacloprid 200 SL	S	0.30 mL/L	1.0	60
VI	Thrips				
	--				
VII	Jassids				
	--				
VIII	Mealy bugs				
26.	Buprofezin 25 SC	NS	1.25 mL/L	1.0	40
27.	Methomyl 40 SP	NS	1.0 g/L	0.02	61 (one application only at pre-flowering stage)
IX	Caterpillars (Helicoverpa armigera and Spodoptera litura)				
	--				
X	Mites				
	--				
XI	Plant Growth Regulators				
28.	Hydrogen cyanamide 50% SL	S	30-40 mL/L	0.05	90-120
29.	Forchlorfenuron (CPPU) 0.1% L	S	1-2 ppm	0.05	45 (for 1 ppm dose) 60 (for 2 ppm dose)

Sr. No.	Chemical recommended for major disease & pest	Nature of chemical	Dose on formulation basis	EU MRL (mg/kg) updated as on 19/07/2010	Pre-harvest Interval (PHI in days)
30.	Gibberellic acid (GA3) Technical	S	100 ppm (Cumulative Usage)	5.00	PHI data not available
31.	1-Naphthyl acetic acid 4.5% L	S	100 ppm	0.05	PHI data not available
32.	Chlormequat chloride 50% SL	S	250 ppm	0.05	PHI data not available
XII	Herbicides				
33.	Paraquat dichloride 24 SL	NS	5 mL/L	0.02	PHI data not available

NS = Non-systemic, S = Systemic

#### Note

- All the doses mentioned above are for high volume sprayers, where normal spray volume is 1000 L/ha. Spray volume can however be changed as per the efficiency of sprayers used. However, the amount of each pesticide based on its active ingredient recommended for 1 ha area on the basis of 1000 L spray solution should be strictly maintained to minimize pesticide residues.
- Recommended PHI will be valid only if two applications of an agrochemical are given per fruiting season at the interval of 7-15 days at recommended dose except in case of Flusilazole.  
  
The PHI of the fungicide Flusilazole pertains to one application by foliar spray only.
- For the chemicals (Sl. No. 30, 31, 32 and 33) having label claim for grapes with CIB&RC, no PHI data available for EU MRL. The farmers need to be cautious about the dose and stage of application as per the need.
- Presently, no insecticides are available with the label claim on grapes with CIB&RC for management of post-harvest berry rots, Thrips, Jassids, Caterpillar and Mites.
- All the recommendations in the Annexure-5 are advisory in nature and therefore the growers are responsible for the compliance to the requirements of their individual respective buyers in respect of chemicals given in Annexure-5.